

Directions for Enfamil LIPIL® Infant Formula

Caution: Carefully follow the instructions of your baby's doctor. The health of your baby may depend on it. It is important to use the directions for the product you are using and the scoop that comes in the can.

22 Cal/fl oz*

From Concentrated Liquid



+

1¼ cup

=



23½ fl oz

1 13 fl oz can concentrate

10½ fl oz (1¼ cup plus 1 Tbsp) water

23½ fl oz infant formula at 22 Cal/fl oz

From Powder



+

¾ cup

=

6 fl oz

3 unpacked level scoops powder

5½ fl oz (¾ cup) water

6 fl oz infant formula at 22 Cal/fl oz

Or

(Use these instructions only if feeding 32 fl oz within 24 hours.)



+

3 cups

=

27 fl oz

1 unpacked level standard household measuring cup

24 fl oz (3 cups) water

27 fl oz infant formula at 22 Cal/fl oz

24 Cal/fl oz*

From Concentrated Liquid



+

1 cup

=



21½ fl oz

1 13 fl oz can concentrate

8½ fl oz (1 cup plus 1 Tbsp) water

21½ fl oz infant formula at 24 Cal/fl oz

From Powder



+

5 fl oz

=

5½ fl oz

3 unpacked level scoops powder

5 fl oz water

5½ fl oz infant formula at 24 Cal/fl oz

Or

(Use these instructions only if feeding 32 fl oz within 24 hours.)



+

3 cups

=

27 fl oz

1 unpacked level standard household measuring cup and 1 unpacked level scoop powder

24 fl oz (3 cups) water

27 fl oz infant formula at 24 Cal/fl oz

Your baby's health depends on carefully following the instructions here and on the product label. Proper hygiene, preparation, dilution, use, and storage are important when preparing infant formula. Powdered infant formulas are not sterile and should **not** be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor. See product label for additional important instructions. **Do not use a microwave oven to warm formulas. Serious burns may result.**

Once prepared, infant formula can spoil quickly. Either immediately feed or immediately cover and store in refrigerator at 35-40°F (2-4°C) for no longer than 24 hours. Do not use prepared formula if it is unrefrigerated for more than a total of 2 hours. Do not freeze prepared formula. After feeding begins, do not refrigerate feeding bottle. You must use within 1 hour or discard.

*Final Cal/fl oz are approximations

Consult your baby's doctor before making feeding changes.

